

Spectral Properties of Some Combinatorial Matrices

EMRAH KILIC¹, GABRIELA N. STĂNICĂ², PANTELIMON STĂNICĂ^{3*}

¹ TOBB Economics and Technology University, Mathematics Department
06560 Sogutozu, Ankara, Turkey; ekilic@etu.edu.tr

² Department of Mathematics. California State University – Monterey Bay
Monterey, CA 93955, USA; gnstanica@gmail.com

³ Department of Applied Mathematics, Naval Postgraduate School
Monterey, CA 93943, USA; pstanica@nps.edu

November 30, 2008

Abstract

In this paper we investigate the spectra and related questions for various combinatorial matrices, generalizing work by Carlitz, Cooper and Kennedy.

1 Introduction

In [8], R. Peele and P. Stănică studied $n \times n$ matrices with the (i, j) entry the binomial coefficient $\binom{i-1}{j-1}$, respectively, $\binom{i-1}{n-j}$ and derived many interesting results on powers of these matrices. In [10], one of us found that the same is true for a much larger class of what he called *netted matrices*, namely matrices with entries satisfying a certain type of recurrence among the entries of all 2×2 cells.

Let R_n be the matrix whose (i, j) entries are $a_{i,j} = \binom{i-1}{n-j}$, which satisfy

$$a_{i,j-1} = a_{i-1,j-1} + a_{i-1,j}. \quad (1)$$

The previous recurrence can be extended for $i \geq 0, j \geq 0$, using the boundary conditions $a_{1,n} = 1, a_{1,j} = 0, j \neq n$. Remark the following consequences of the boundary conditions and recurrence (1): $a_{i,j} = 0$ for $i + j \leq n$, and $a_{i,n+1} = 0, 1 \leq i \leq n$.

The matrix R_n was firstly studied by Carlitz [2] who gave explicit forms for the eigenvalues of R_n . Let $f_{n+1}(x) = \det(xI - R_n)$ be the characteristic polynomial of R_n . Thus

$$f_n(x) = \sum_{r=0}^{n+1} (-1)^{r(r+1)/2} \binom{n}{r}_F x^{n-r}$$

*Also Associated to the Institute of Mathematics of Romanian Academy, Bucharest, Romania

| Report Documentation Page | | | | Form Approved OMB No. 0704-0188 | |
|--|------------------------------------|-------------------------------------|---|---|---------------------------------|
| Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. | | | | | |
| 1. REPORT DATE 30 NOV 2008 | | 2. REPORT TYPE | | 3. DATES COVERED 00-00-2008 to 00-00-2008 | |
| 4. TITLE AND SUBTITLE Spectral Properties of Some Combinatorial Matrices | | | | 5a. CONTRACT NUMBER | |
| | | | | 5b. GRANT NUMBER | |
| | | | | 5c. PROGRAM ELEMENT NUMBER | |
| 6. AUTHOR(S) | | | | 5d. PROJECT NUMBER | |
| | | | | 5e. TASK NUMBER | |
| | | | | 5f. WORK UNIT NUMBER | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Postgraduate School, Department of Applied Mathematics, Monterey, CA, 93943 | | | | 8. PERFORMING ORGANIZATION REPORT NUMBER | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) | | | | 10. SPONSOR/MONITOR'S ACRONYM(S) | |
| | | | | 11. SPONSOR/MONITOR'S REPORT NUMBER(S) | |
| 12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited | | | | | |
| 13. SUPPLEMENTARY NOTES Congressus Numerantium Vol. 201, Proc. International Conf. Fibonacci Numbers & Appl., (F. Luca, P. Stanica, eds.) (2010), 223-236. | | | | | |
| 14. ABSTRACT In this paper we investigate the spectra and related questions for various combinatorial matrices, generalizing work by Carlitz, Cooper and Kennedy. | | | | | |
| 15. SUBJECT TERMS | | | | | |
| 16. SECURITY CLASSIFICATION OF: | | | 17. LIMITATION OF ABSTRACT Same as Report (SAR) | 18. NUMBER OF PAGES 12 | 19a. NAME OF RESPONSIBLE PERSON |
| a. REPORT unclassified | b. ABSTRACT unclassified | c. THIS PAGE unclassified | | | |

where $\binom{n}{r}_F$ denote the Fibonomial coefficient, defined (for $n \geq r > 0$) by

$$\binom{n}{r}_F = \frac{F_1 F_2 \dots F_n}{(F_1 F_2 \dots F_r)(F_1 F_2 \dots F_{n-r})}$$

with $\binom{n}{n}_F = \binom{n}{0}_F = 1$. Carlitz showed that

$$f_n(x) = \prod_{j=0}^{n-1} (x - \phi^j \bar{\phi}^{n-j})$$

where $\phi, \bar{\phi} = (1 \pm \sqrt{5})/2$. Thus the eigenvalues of R_n are $\phi^n, \phi^{n-1}\bar{\phi}, \dots, \phi\bar{\phi}^{n-1}, \bar{\phi}^n$. We shall give another proof of this result in the next section.

In [8] it was proved that the entries of the power R_n^e satisfy the recurrence

$$F_{e-1}a_{i,j}^{(e)} = F_e a_{i-1,j}^{(e)} + F_{e+1}a_{i-1,j-1}^{(e)} - F_e a_{i,j-1}^{(e)}, \quad (2)$$

where F_e is the Fibonacci sequence. Closed forms for *all* entries of R_n^e were not found, but several results concerning the generating functions of rows and columns were obtained (see [8, 10]). For instance, the entries in the first row and column of R_n^e are

$$\begin{aligned} a_{1,j}^{(e)} &= \binom{n-1}{j-1} F_{e-1}^{n-j} F_e^{j-1} \\ a_{i,1}^{(e)} &= F_{e-1}^{n-i} F_e^{i-1}. \end{aligned} \quad (3)$$

Further, the generating function for the (i, j) -th entry of the e -th power of a generalization of R_n , namely

$$Q_n(a, b) = \left(a^{i+j-n-1} b^{n-j} \binom{i-1}{n-j} \right)_{1 \leq i, j \leq n}$$

is

$$B_n^{(e)}(x, y) = \frac{(U_{e-1} + U_e y)(b U_{e-1} + y U_e)^{n-1}}{U_{e-1} + U_e y - x(U_e + U_{e+1} y)}.$$

Certainly, $Q_n(1, 1) = R_n$. As an example,

$$Q_6(a, b) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & b & a \\ 0 & 0 & 0 & b^2 & 2ab & a^2 \\ 0 & 0 & b^3 & 3ab^2 & 3a^2b & a^3 \\ 0 & b^4 & 4ab^3 & 6a^2b^2 & 4a^3b & a^4 \\ b^5 & 5ab^4 & 10a^2b^3 & 10a^3b^2 & 5a^4b & a^5 \end{pmatrix}.$$

Regarding this generalization, in [3], the authors gave the characteristic polynomial of $Q_n(a, b)$ and the trace of k th power of $Q_n(a, b)$, that is, $\text{tr}(Q_n^k(a, b))$, by using the method of Carlitz [2].

Lemma 1. Let $g_n(x)$ denote the characteristic polynomial of $Q_n(a, b)$. Thus

$$g_n(x) = \sum_{i=0}^n (-1)^{i(i+1)/2} b^{i(i-1)/2} \binom{n}{i}_U x^{n-i} \quad (4)$$

and

$$\text{tr}(Q_n^k(a, b)) = \frac{U_{kn}}{U_k}, \quad (5)$$

where $\binom{n}{i}_U$ stands for the generalized Fibonomial coefficient, defined by

$$\binom{n}{r}_U = \frac{U_1 U_2 \dots U_n}{(U_1 U_2 \dots U_r)(U_1 U_2 \dots U_{n-r})},$$

for $n \geq i > 0$, where $\binom{n}{n}_U = \binom{n}{0}_U = 1$.

One of us extended in [10] some of these results to netted matrices, say $R_n^{\alpha, \beta, \gamma, \delta}$, whose entries $a_{i,j}$, $i \geq 0$, $j \geq 0$ satisfy (for $i \geq 1$, $j \geq 1$)

$$\delta a_{i,j} = \alpha a_{i-1,j} + \beta a_{i-1,j-1} + \gamma a_{i,j-1}, \quad (6)$$

with the boundary conditions

$$\beta a_{i,0} + \gamma a_{i+1,0} = 0, \text{ for all } 1 \leq i \leq n-1 \quad (7)$$

$$\delta a_{i+1,n+1} - \alpha a_{i,n+1} = 0, \text{ for all } 1 \leq i \leq n-1. \quad (8)$$

It was shown in [10] that the entries of any power of such a matrix will satisfy a similar recurrence, and generating functions for these powers were found. In the case of the Fibonacci sequence, the generating function of the i -th row of R_n^e is

$$r_i^{(e)}(x) = \sum_{j \geq 1} a_{i,j}^{(e)} x^{j-1} = (F_e + F_{e+1}x)^{i-1} (F_{e-1} + F_e x)^{n-i}$$

and, if $e > 1$, the generating function for the j -th column of R_n^e is

$$c_j^{(e)}(x) = \left(\frac{F_{e+1}x - F_e}{F_{e-1} - F_e x} \right)^{j-1} \frac{F_{e-1}^n}{F_{e-1} - F_e x} \left[1 + \sum_{s=1}^{j-1} \binom{n}{s} \left(\frac{F_e(F_{e-1} - F_e x)}{F_{e-1}(F_{e+1}x - F_e)} \right)^s \right].$$

where $c_j^{(1)}(x) = (1-x)^{j-1-n} x^{n-j}$.

The matrix $Q_n(a, b)$ was introduced as a generalization of the Fibonacci matrix $Q = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$ which has the property that $Q^n \cdot \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} F_{n-1} \\ F_n \end{pmatrix}$. The matrix $Q_n(a, b)$ displays a similar property on the sequence $U_{n+1} = aU_n + bU_{n-1}$, that is, any e -th power of $Q_n(a, b)$ multiplied by a fixed vector gives an n -tuple of consecutive terms of the sequence $U = (U_k)_k$. We refer to [10, 3] for more details.

2 Spectra of R_n and $Q_n(a, b)$

In [8], the authors proposed a conjecture on the eigenvalues, which was proven independently in [1] and the unpublished manuscript [11]. In this section we give the proof of the conjecture from [11] and we find the eigenvectors of R_n . In [8, 10], it was shown that the inverse of R_n is the matrix

$$R_n^{-1} = \left((-1)^{n+i+j+1} \binom{n-i}{j-1} \right)_{1 \leq i, j \leq n},$$

and, in general, the inverse of $Q_n(a, b)$ is

$$Q_n^{-1}(a, b) = \left((-1)^{n+i+j+1} a^{n+1-i-j} b^{i-n} \binom{n-i}{j-1} \right)_{1 \leq i, j \leq n}. \quad (9)$$

We define K_n to be the matrix with (i, j) -entry $\delta_{i, n-j+1}$ (the Kronecker symbol), that is, K_n is a permutation matrix having 1 on the secondary diagonal and 0 elsewhere.

Theorem 2. *Let $\phi = \frac{1+\sqrt{5}}{2}$, $\bar{\phi} = \frac{1-\sqrt{5}}{2}$ be the golden section and its conjugate. The eigenvalues of R_n are:*

1. $\{(-1)^{k+i} \phi^{2i-1}, (-1)^{k+i} \bar{\phi}^{2i-1}\}_{i=1, \dots, k}$, if $n = 2k$.
 2. $\{(-1)^k\} \cup \{(-1)^{k+i} \phi^{2i}, (-1)^{k+i} \bar{\phi}^{2i}\}_{i=1, \dots, k}$, if $n = 2k + 1$.
- Equivalently, the eigenvalues of R_n are $\phi^n, \phi^{n-1} \bar{\phi}, \dots, \phi \bar{\phi}^{n-1}, \bar{\phi}^n$.*

Proof. First, we show that R_n is a permutation matrix away from L_n , namely

$$R_n \cdot K_n = L_n \iff R_n = L_n \cdot K_n. \quad (10)$$

It is a trivial matter to prove $K_n^2 = I_n$, which will give the equivalence. It suffices to show the second identity, which follows easily, since an entry in $L_n \cdot K_n$ is

$$\sum_{k=1}^n \binom{i-1}{k-1} \delta_{k, n-j+1} = \binom{i-1}{n-j}.$$

Now, denote by A_n , the matrix obtained by taking absolute values of entries of R_n^{-1} . We show that R_n has the same characteristic polynomial (eigenvalues) as A_n , namely we prove their similarity,

$$K_n \cdot R_n \cdot K_n = A_n. \quad (11)$$

Since $K_n \cdot R_n \cdot K_n = K_n \cdot L_n$ (by (10)), to show (11) it suffices to prove that $K_n \cdot L_n = \left(\binom{n-i}{j-1} \right)_{i, j}$. Therefore, we need

$$\sum_{k=1}^n \delta_{i, n-k+1} \binom{k-1}{j-1} = \binom{n-i}{j-1},$$

which is certainly true.

We use a result of [7] to show that A_n in turn is similar to D_n , the diagonal matrix whose diagonal entries are the elements in the eigenvalues set listed in decreasing order according to size of the absolute value. For instance, for $n = 4$,

$$D_4 = \begin{pmatrix} \alpha^3 & 0 & 0 & 0 \\ 0 & -\alpha & 0 & 0 \\ 0 & 0 & -\beta & 0 \\ 0 & 0 & 0 & \beta^3 \end{pmatrix}$$

Our claim will be proved if one can show that A_n is similar to D_n , and this will follow from [7]. We sketch here that argument for the convenience of the reader. Define an array $b_{n,m}$ by

$$\begin{aligned} b_{n,0} &= 1 \text{ for all } n \geq 0 \\ b_{n,m} &= 0 \text{ for all } m > n \\ b_{n,m} &= b_{n-1,m-1} (-1)^m \frac{F_n}{F_m} \text{ for all } m \leq n \end{aligned}$$

Certainly, $|b_{n,m}| = \frac{F_n F_{n-1} \cdots F_{n-m+1}}{F_m \cdots F_1}$. Let $C_n = (c_{i,j})_{i,j}$, where

$$\begin{cases} c_{i,i+1} = 1 & \text{if } i = 1, \dots, n-1 \\ c_{n,j} = -b_{n,n+1-j} & \text{if } j = 1, \dots, n \\ c_{i,j} = 0 & \text{otherwise.} \end{cases}$$

We observe that, in fact, C_n is the companion matrix of the polynomial with coefficients $b_{n,n+1-j}$. Let X_n be the matrix with entries $\binom{n-i}{j-1} F_{i-2}^{j-1} F_{i-1}^{n-j}$. It turns out that the eigenvector matrix E_n of A_n , with columns vectors listed in decreasing order of absolute value of the corresponding eigenvalues, normalized so that the last row is made up of all 1's, satisfies

$$X_n E_n = V_n,$$

where V_n is the Vandermonde matrix, which is the eigenvector matrix of C_n with eigenvectors listed in decreasing order of the absolute values of the corresponding eigenvalues. Also,

$$X_n A_n X_n^{-1} = C_n \text{ and } E_n^{-1} A_n E_n = D_n.$$

Our theorem follows. ■

Easily we deduce

Corollary 3. *The eigenvectors matrix of R_n , say W_n , with eigenvectors listed in decreasing order of the absolute values of the corresponding eigenvalues, is*

$$W_n = K_n E_n = K_n X_n^{-1} V_n.$$

Proof. We showed that $K_n R_n K_n = A_n$ and $E_n^{-1} A_n E_n = D_n$. It follows that $(E_n^{-1} K_n) R_n (K_n E_n) = D_n$, which together with $X_n E_n = V_n$, proves the corollary. ■

Example 4. For $n = 4$, the eigenvectors matrix is

$$W_4 = \begin{pmatrix} -\alpha^3 & \alpha & \beta & -\beta^3 \\ \alpha^2 & -\frac{1}{3}\beta & -\frac{1}{3}\alpha & \beta^2 \\ -\alpha & -\frac{1}{3}\alpha^2 & -\frac{1}{3}\beta^2 & -\beta \\ 1 & 1 & 1 & 1 \end{pmatrix}$$

We present below the first few characteristic polynomials of $Q_n(a, b)$ (for $1 \leq n \leq 7$) and we ask whether one can find their roots as in Theorem 2:

$$\begin{aligned} & x^2 - ax - b; \\ & -(b+x)(-xa^2 + b^2 + x^2 - 2bx); \\ & (b^3 - axb - x^2)(xa^3 + 3bxa + b^3 - x^2); \\ & (b^2 - x)(-xa^4 - 4bxa^2 + b^4 + x^2 - 2b^2x)(b^4 + 2xb^2 + a^2xb + x^2); \\ & -(b^5 - 3axb^2 - a^3xb - x^2)(b^5 + axb^2 - x^2)(xa^5 + 5bxa^3 + 5b^2xa + b^5 - x^2). \end{aligned}$$

The argument that proves the next theorem is similar to the proof of Theorem 2, and so, we omit it. One can also find two different ways to see its proof in [3] and [4].

Theorem 5. Let $\lambda = \frac{a+\sqrt{a^2+4b}}{2}$, $\bar{\lambda} = \frac{a-\sqrt{a^2+4b}}{2}$ be the roots of the polynomial $x^2 - ax - b$. The eigenvalues of $Q_n(a, b)$ are:

1. $\{(-b)^{k-i} \lambda^{2i-1}, (-b)^{k-i} \bar{\lambda}^{2i-1}\}_{i=1, \dots, k}$, if $n = 2k$.
2. $\{(-b)^k\} \cup \{(-b)^{k-i} \lambda^{2i}, (-b)^{k-i} \bar{\lambda}^{2i}\}_{i=1, \dots, k}$, if $n = 2k+1$. Equivalently, the eigenvalues of $Q_n(a, b)$ can also be written in the following compact form $\lambda^n, \lambda^{n-1}\bar{\lambda}, \dots, \lambda\bar{\lambda}^{n-1}, \bar{\lambda}^n$.

Since $R_n, Q_n(a, b)$ are column justified triangular matrices, the determinants of our matrices are easy to compute, or we can use the previous result.

Corollary 6. We have $\det(R_n) = (-1)^{\lfloor n/2 \rfloor}$ and $\det(Q_n(a, b)) = (-1)^{\lfloor n/2 \rfloor} b^{n(n-1)/2}$.

3 Arithmetic progressions and spectra of other combinatorial matrices

Let the sequences $\{u_n\}, \{v_n\}$ be defined by

$$\begin{aligned} u_n &= au_{n-1} + bu_{n-2} \\ v_n &= av_{n-1} + bv_{n-2}, \end{aligned}$$

for $n > 1$, where $u_0 = 0, u_1 = 1$, and $v_0 = 2, v_1 = a$, respectively. Let α, β be the roots of the associated equation $x^2 - ax - b = 0$. The next lemma appears in [6].

Lemma 7. For $k \geq 1$ and $n > 1$,

$$\begin{aligned} u_{kn} &= v_k u_{k(n-1)} + (-1)^{k+1} b^k u_{k(n-2)} \\ v_{kn} &= v_k v_{k(n-1)} + (-1)^{k+1} b^k v_{k(n-2)}. \end{aligned} \tag{12}$$

Using the sequence v_k , we define the $n \times n$ matrix $H_n(v_k, b^k)$ as follows:

$$H_n(v_k, b^k) = \left(v_k^{i+j-n-1} \left(-(-b)^k \right)^{n-j} \binom{i-1}{n-j} \right)_{1 \leq i, j \leq n}.$$

Observe that $H_n(v_1, b^1) = Q_n(a, b)$. As an example,

$$H_6(v_k, b^k) = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & -(-b)^k & v_k \\ 0 & 0 & 0 & b^{2k} & -2(-b)^k v_k & v_k^2 \\ 0 & 0 & -(-b)^{3k} & 3b^{2k} v_k & -3(-b)^k v_k^2 & v_k^3 \\ 0 & b^{4k} & -4(-b)^{3k} v_k & 6b^{2k} v_k^2 & -4(-b)^k v_k^3 & v_k^4 \\ -(-b)^{5k} & 5b^{4k} v_k & -10(-b)^{3k} v_k^2 & 10b^{2k} v_k^3 & -5(-b)^k v_k^4 & v_k^5 \end{pmatrix}.$$

As in equation (9), we can easily find the inverse of the matrix H_n , namely

$$H_n^{-1}(v_k, b^k) = \left((-1)^{j+1} (-b)^{-k(n-i)} v^{n+1-i-j} \binom{n-i}{j-1} \right)_{i,j}.$$

It is well known that for $n \geq -1$,

$$u_{n+1} = \sum_r \binom{r}{n-r} a^{2r-n} b^{n-r}. \quad (13)$$

We generalize this identity next.

Lemma 8. For $k > 0$ and $n \geq -1$,

$$\frac{u_{k(n+1)}}{u_k} = \sum_r \binom{r}{n-r} v_k^{2r-n} \left(-(-b)^k \right)^{n-r}.$$

Proof. (Induction on n) If $n = 1$, then equations (12) and (13) give the identity. Suppose that the result is true for $n - 1$ and n . Then

$$\begin{aligned} u_{k(n+1)} &= v_k u_{kn} + (-b)^{k+1} u_{k(n-1)} \\ &= v_k \sum_r \binom{r}{n-1-r} v_k^{2r-n+1} \left(-(-b)^k \right)^{n-1-r} \\ &\quad - (-b)^k \sum_r \binom{r}{n-2-r} v_k^{2r-n+2} \left(-(-b)^k \right)^{n-2-r} \\ &= \sum_r \left[\binom{r}{n-1-r} + \binom{r}{n-2-r} \right] v_k^{2r-n+2} \left(-(-b)^k \right)^{n-1-r} \\ &= \sum_r \binom{r+1}{n-1-r} v_k^{2r-n+2} \left(-(-b)^k \right)^{n-1-r} \\ &= \sum_r \binom{r}{n-r} v_k^{2r-n} \left(-(-b)^k \right)^{n-r}. \end{aligned}$$

■

Lemma 9. Let $k, m > 0$ and $0 \leq r \leq n$. Then

$$\begin{aligned} & \left(\frac{u_{km}x - (-b)^m u_{(k-1)m}}{u_m} \right)^r \left(\frac{u_{(k+1)m}x - (-b)^m u_{km}}{u_m} \right)^{n-r} \\ &= \sum_{r_1, r_2, \dots, r_k} \binom{n-r}{r_1} \binom{n-r_1}{r_2} \dots \binom{n-r_{k-1}}{r_k} \\ & \quad \times v_m^{kn-r-2r_1-\dots-2r_{k-1}-r_k} (-(-b)^m)^{r_1+r_2+\dots+r_k} x^{n-r_k}. \end{aligned} \quad (14)$$

Proof. (Induction on k) For $k = 1$, the proof follows from $u_{2n} = u_n v_n$ and

$$x^r (v_m x - (-b)^m)^{n-r} = \sum_s \binom{n-r}{s} v_m^{n-r-s} (-(-b)^m)^s x^{n-s}$$

(see [3]). We assume that the equality holds for some positive integer k . If we replace $(v_m - (-b)^m x^{-1})$ by x and then multiply by x^n , the left side of this equation becomes

$$\left(\frac{u_{(k+1)m}x - (-b)^m u_{km}}{u_m} \right)^r \left(\frac{u_{(k+2)m}x - (-b)^m u_{(k+1)m}}{u_m} \right)^{n-r}.$$

After some simplifications, if we expand the right side of this equation, then we get

$$\begin{aligned} & \sum_{r_1, r_2, \dots, r_{k+1}} \binom{n-r}{r_1} \binom{n-r_1}{r_2} \dots \binom{n-r_k}{r_{k+1}} \\ & \times v_m^{k(n+1)-r-2r_1-\dots-2r_k-r_{k+1}} (-(-b)^m)^{r_1+r_2+\dots+r_{k+1}} x^{n-r_{k+1}}. \end{aligned}$$

The proof is complete. ■

Lemma 10. For all $m > 0$,

$$\text{tr} \left(H_n^m (v_k, b^k) \right) = \frac{u_{knm}}{u_k}.$$

Proof. If we multiply both sides of the equation (14) by x^r and summing over r , we get

$$\begin{aligned} & \sum_{r=0}^n \left(\frac{u_{km}x - (-b)^m u_{(k-1)m}}{u_m} \right)^r \left(\frac{u_{(k+1)m}x - (-b)^m u_{km}}{u_m} \right)^{n-r} x^r \\ &= \sum_{r_1, r_2, \dots, r_k} \binom{n-r}{r_1} \binom{n-r_1}{r_2} \dots \binom{n-r_{k-1}}{r_k} \\ & \quad \times v_m^{kn-r-2r_1-\dots-2r_{k-1}-r_k} (-(-b)^m)^{r_1+r_2+\dots+r_k} x^{n+r-r_k}, \end{aligned}$$

where the coefficient of x^n on the right side is $\text{tr} (H_n^m (v_k, b^k))$. The coefficient of x^n on the left side of this equation without its denominator is

$$\begin{aligned} & \sum_{r+s+t=n} \binom{r}{s} \binom{n-r}{t} u_{km}^s (-(-b)^m u_{(k-1)m})^{r-s} u_{(k+1)m}^t (-(-b)^m u_{km})^s \\ &= \sum_{r+s \leq n} \binom{r}{s} \binom{n-r}{s} u_{km}^s (-(-b)^m u_{(k-1)m})^{r-s} u_{(k+1)m}^{n-r-s} (-(-b)^m u_{km})^s. \end{aligned}$$

For easy writing, denote this last expression by c_n . Then

$$\begin{aligned}
& \sum_{n=0}^{\infty} c_n x^n \\
&= \sum_{r,s=0}^{\infty} \binom{r}{s} (-(-b)^m)^r u_{(k-1)m}^{r-s} u_{km}^{2s} x^{r+s} \sum_{n=r+s}^{\infty} \binom{n-r}{s} (u_{(k+1)m} x)^{n-r-s} \\
&= \sum_{r,s=0}^{\infty} \binom{r}{s} (-(-b)^m)^r u_{(k-1)m}^{r-s} u_{km}^{2s} x^{r+s} (1 - u_{(k+1)m} x)^{-s-1} \\
&= \sum_{s=0}^{\infty} (-(-b)^m)^s u_{km}^{2s} x^{2s} (1 - u_{(k+1)m} x)^{-s-1} \sum_{r \geq s} \binom{r}{s} ((-(-b)^m) u_{(k-1)m} x)^{r-s} \\
&= \sum_{s=0}^{\infty} (-(-b)^m)^s u_{km}^{2s} x^{2s} (1 - u_{(k+1)m} x)^{-s-1} (1 + (-b)^m u_{(k-1)m} x)^{-s-1} \\
&= \frac{1}{(1 - u_{(k+1)m} x) (1 + (-b)^m u_{(k-1)m} x)} \frac{1}{1 - \frac{(-(-b)^m)^s u_{km}^{2s} x^2}{(1 - u_{(k+1)m} x) (1 + (-b)^m u_{(k-1)m} x)}} \\
&= \frac{1}{(1 - u_{(k+1)m} x) (1 + (-b)^m u_{(k-1)m} x) + (-b)^m u_{km}^{2s} x^2}. \tag{15}
\end{aligned}$$

Here by the Binet formula for $\{u_n\}$, we note that for all integers $k > 0$,

$$\frac{u_{kn} - (-b)^k u_{k(n-2)}}{u_k} = v_{k(n-1)} \tag{16}$$

and

$$-\left(\frac{u_{kn} u_{k(n-2)} - u_{k(n-1)}^2}{u_k^2} \right) = (-b)^{k(n-2)}. \tag{17}$$

Using (16) and (17), we write the right side of the equation (15) as

$$\begin{aligned}
& \frac{1}{(1 - u_{(k+1)m} x) (1 + (-b)^m u_{(k-1)m} x) + (-b)^m u_{km}^{2s} x^2} \\
&= \frac{1}{1 - v_{mk} x + (-b)^{mk} x^2}.
\end{aligned}$$

Thus, $c_n = \frac{u_{knm}}{u_k}$, and so, $\text{tr}(H_n^m(v_k, b^k)) = \frac{u_{knm}}{u_k}$. The proof is complete. ■

Theorem 11. *The eigenvalues of $H_n(v_k, b^k)$ are*

$$\alpha^{kn}, \alpha^{k(n-1)} \beta^k, \dots, \alpha^k \beta^{k(n-1)}, \beta^{kn}.$$

Proof. Let $f_n(x) = \det(xI - H_n(v_k, b^k))$ and $\lambda_0, \lambda_1, \dots, \lambda_{n-1}$ denote the eigenvalues of $H_n(v_k, b^k)$. Then by Lemma 10,

$$\begin{aligned}
\frac{f'_n(x)}{f_n(x)} &= \sum_{j=0}^{n-1} \frac{1}{x - \lambda_j} = \sum_{m=0}^{n-1} x^{-m-1} \sum_{j=0}^m \lambda_j^m \\
&= \sum_{k=0}^{\infty} x^{-k-1} \text{tr} \left(H_{n-1}^m(v_k, b^k) \right) \\
&= \sum_{m=0}^{\infty} x^{-m-1} \frac{u_{k(n-1)m}}{u_k} \\
&= \sum_{m=0}^{\infty} x^{-m-1} \sum_{j=0}^{n-1} \alpha^{jkm} \beta^{k(n-j-1)m} \\
&= \sum_{j=0}^{n-1} \frac{1}{x - \alpha^{jk} \beta^{(n-j-1)k}}.
\end{aligned}$$

Thus

$$f_n(x) = \prod_{j=0}^{n-1} \left(x - \alpha^{jk} \beta^{(n-j-1)k} \right)$$

and so the eigenvalues of $H_n(v_k, b^k)$ are

$$\alpha^{k(n-1)}, \alpha^{k(n-2)} \beta^k, \dots, \alpha^k \beta^{k(n-2)}, \beta^{k(n-1)}.$$

■

Next, we give an expression of the characteristic polynomial of H_n in terms of Gaussian binomials. Fix k and let $r_n := u_{kn}$. Define the k -Fibonomial coefficients by

$$\left\{ \begin{matrix} n \\ m \end{matrix} \right\}_{u,k} = \frac{r_1 r_2 \dots r_n}{(r_1 r_2 \dots r_m) (r_1 r_2 \dots r_{n-m})},$$

where $\left\{ \begin{matrix} n \\ n \end{matrix} \right\}_{u,k} = \left\{ \begin{matrix} n \\ 0 \end{matrix} \right\}_{u,k} = 1$.

Clearly, when $k = 1$, then $\left\{ \begin{matrix} n \\ m \end{matrix} \right\}_{u,k}$ is reduced to the Fibonomial coefficient $\left(\begin{matrix} n \\ m \end{matrix} \right)_u$.

Theorem 12.

$$f_n(x) = \prod_{j=0}^{n-1} \left(x - \alpha^{jk} \beta^{(n-j-1)k} \right) = \sum_{i=0}^n (-1)^i (-b)^{ki(i-1)/2} \left\{ \begin{matrix} n \\ i \end{matrix} \right\}_{u,k} x^{n-i}.$$

Proof. Here we can use the following familiar identity

$$\prod_{j=0}^{n-1} (1 - q^j x) = \sum_{i=0}^n (-1)^i q^{i(i-1)/2} \left[\begin{matrix} n \\ i \end{matrix} \right]_q x^i, \quad (18)$$

where

$$\begin{bmatrix} n \\ i \end{bmatrix}_q = \frac{(1 - q^n) \dots (1 - q^{n-i+1})}{(1 - q^i) \dots (1 - q)},$$

are the usual q -binomial coefficients (Gaussian binomials).

If we replace q by $(\beta/\alpha)^k$ we find that for all positive fixed integers k ,

$$\begin{bmatrix} n \\ i \end{bmatrix}_q \rightarrow \alpha^{ki(i-n)} \left\{ \begin{matrix} n \\ i \end{matrix} \right\}_{u,k}.$$

Thus (18) becomes

$$\prod_{j=0}^{n-1} \left(1 - (\alpha^{-1}\beta)^{kj} x \right) = \sum_{i=0}^n (-1)^i \beta^{ki(i-1)/2} \alpha^{ki(i+1)/2 - nki} \left\{ \begin{matrix} n \\ i \end{matrix} \right\}_{u,k} x^i.$$

Replacing x by $\alpha^{k(n-1)}x$, we get

$$\begin{aligned} \prod_{j=0}^{n-1} \left(1 - \alpha^{(n-j-1)k} \beta^{kj} x \right) &= \sum_{i=0}^n (-1)^i (\alpha\beta)^{ki(i-1)/2} \left\{ \begin{matrix} n \\ i \end{matrix} \right\}_{u,k} x^i \\ &= \sum_{i=0}^n (-1)^i (-b)^{ki(i-1)/2} \left\{ \begin{matrix} n \\ i \end{matrix} \right\}_{u,k} x^i. \end{aligned}$$

Finally replacing x by x^{-1} gives us

$$\prod_{j=0}^{n-1} \left(1 - \alpha^{(n-j-1)k} \beta^{kj} x \right) = \sum_{i=0}^n (-1)^i (-b)^{ki(i-1)/2} \left\{ \begin{matrix} n \\ i \end{matrix} \right\}_{u,k} x^{n-i}.$$

■

Acknowledgements. The authors would like to thank the referee for constructive comments, which improved the presentation of the paper. The third author acknowledges NPS Foundation and NPS-RIP funding support from Naval Postgraduate School.

References

- [1] D. Callan and H. Prodinger, *An involutory matrix of eigenvectors*, Fibonacci Quart. **41:2** (2003), 105–107.
- [2] L. Carlitz, *The characteristic polynomial of a certain matrix of binomial coefficients*, Fibonacci Quart. **3** (1965), 81–89.
- [3] C. Cooper and R. Kennedy, *Proof of a result by Jarden by generalizing a proof by Carlitz*, Fibonacci Quart. **33:4** (1995), 304–310.
- [4] P. Duvall and T. Vaughan, *Pell polynomials and a conjecture of Mahon and Horadam*, Fibonacci Quart. **26:4** (1988), 344–353.

- [5] R.L. Graham, D.E. Knuth, O. Patashnik, *Concrete Mathematics*, Adison-Wesley Publishing Company, 1989.
- [6] E. Kilic and P. Stănică, *Factorizations of binary polynomial recurrences by matrix methods*, submitted.
- [7] E. Liverance and J. Pitsenberger, *Diagonalization of the Binomial Matrix*, Fibonacci Quart. **34** (1996), 55–66.
- [8] R. Peele and P. Stănică, *Matrix Powers of Column-Justified Pascal Triangles and Fibonacci Sequences*, Fibonacci Quart. **40** (2002), 146–152.
- [9] M. Petkovsek, H. Wilf, D. Zeilberger, *A = B*, A.K. Peters, Ltd., Wellesley, Massachusetts, 1997.
- [10] P. Stănică, *Netted Matrices*, Int. J. Math. Math. Sci. **39** (2003), 2507–2518.
- [11] P. Stănică and R. Peele, *Spectral Properties of a Binomial Matrix*, unpublished manuscript, <http://arxiv.org/abs/math/0011111>, 2000.
- [12] N. Strauss, *Jordan Form of a Binomial Coefficient matrix over \mathbb{Z}_p* , Linear Alg. Appl. **90** (1987), 60–72.